

"It doesn't matter what other's are doing, it matters what you are doing"

# CSIR NET – Life Science





## Unit 1: Biochemistry

14

Carbohydrate Metabolism



Order Online and Get  
Free Delivery Across India





## Points to be covered in this Lecture



Pasteur Effect



Warburg Effect



Linkage Step



TCA Cycle



Gluconeogenesis



Pentose Phosphate Pathway



Glycogen Metabolism



### Pasteur Effect:

- Presence of oxygen decreases the rate of glycolysis. (fermentation)
- More ATP in presence of oxygen inhibits glycolysis

↓  
mitochondria




↑ ATP —→ PFK-1

↑ ATP —→ Pyruvate Kinase



### ✓ Crabtree Effect:


- High glucose concentrations lead to a preference for glycolysis even in the presence of adequate oxygen.
- More ethanol is produced from glucose instead of entry of pyruvate into mitochondria
- Observed in yeast under high external glucose concentration






## Warburg effect in cancerous cells:

- The enhanced conversion of glucose to lactate observed in tumor cells,
- Even in the presence of normal levels of oxygen





CITRIC ACID CYCLE








## Linkage Step

Conversion of pyruvate into acetyl CoA (Energy Rich molecules)

$3c$                        $2c$





The pyruvate dehydrogenase multienzyme complex (PDC) consists of **3 enzymes**:

- ✓ 1. Pyruvate dehydrogenase ( $E_1$ ) — **TPP**
- ✓ 2. Dihydrolipoyl transacetylase ( $E_2$ ) — **Lipoamide, co-enz A**
- ✓ 3. Dihydrolipoyl dehydrogenase ( $E_3$ ) — **FAD,  $NAD^+$**

The five coenzyme required by this complex are

**$E_1$ : TPP**

**$E_2$ : lipoamide and coenzyme A**

**$E_3$ : FAD and NAD**




**Arsenic compounds sequester lipoamide**





## Regulation of PDH complex

### 1. Product inhibition by NADH and acetyl-CoA



### 2. Covalent modification of the pyruvate dehydrogenase (E<sub>1</sub>) subunit

- Phosphorylated form: Inactive
- Dephosphorylated form: **Active**







## CITRIC ACID CYCLE

Krebs cycle or tricarboxylic acid (TCA)

**Location:** mitochondrial matrix in eukaryotes, cytosol of prokaryotes


Amphibolic in nature

ATP = ↓    NADH = ↓    : Catabolic cycle  
ATP = ↑    NADH = ↑    : Anabolic cycle





## Step 1: Formation of citrate (Tricarboxylic acid)




Citrate inhibits phosphofructokinase I (glycolysis) and activates acetyl CoA carboxylase (fatty acid synthesis).

Inhibitor: citrate, ATP (the  $K_m$  for acetyl CoA is raised as the level of ATP rises).







## Step 2: Isomerization of citrate to iso-citrate





## Step 3: Formation of $\alpha$ -ketoglutarate by Isocitrate dehydrogenase (ICD)




Regulatory step

Inhibitor: NADH, ATP

Activator: ADP

• Irreversible step

## Step 4: Conversion of $\alpha$ -ketoglutarate to succinyl CoA by $\alpha$ -KGDH



• Oxidative Decarboxylation

• Irreversible

$E_1$  : TPP

$E_2$  : Lipoamide, Co-enz A

$E_3$  : FAD,  $\text{NAD}^+$

$\alpha$ -Ketoglutarate TPP, FAD, lipoate

succinyl CoA (4C)

• 3 enzymes ( $E_1, E_2, E_3$ )


• 5 co-enzymes

• Energy Rich molecule

Inhibitor: NADH and succinyl CoA.



## Step 5: Formation of succinate by succinyl-CoA synthetase




Mammals: GTP  
plant and bacteria: ATP





Step 6: Conversion of succinate to fumarate by SDH: *oxidation*


Step 7: Formation of malate by fumarase: *Hydration*



- Inner membrane of mitochondria
- $\text{FAD} \longrightarrow \text{FADH}_2$
- Inhibitor: malonate



## Step 8: Conversion of malate to oxaloacetate by MDH




$$\begin{aligned}
 \Delta G_{\text{overall}}^\circ &= \Delta G_1^\circ + \Delta G_2^\circ \\
 &= (+7.1) + (-7.8) \\
 &= 7.1 - 7.8 \\
 &= -0.7 \text{ Kcal/mole.}
 \end{aligned}$$




## Requirement of $O_2$ by TCA cycle:

### Indirect Requirement for Oxygen:

- Without oxygen, the ETC cannot operate
- $NAD^+$  and FAD regeneration cannot be observed
- TCA cycle would not function



$O_2$  absent






## Summary of TCA cycle:

$$\underline{1 \text{ Pyruvate}} = 12.5 \text{ ATP}$$

The events of Krebs cycle may be summarized as



1 Acetyl CoA  $\Rightarrow$  10 ATP equivalent energy


## Energetics of TCA cycle:

$$3 \text{ NADH} \times 2.5 \text{ ATP} = 7.5 \text{ ATP}$$

$$1 \text{ FADH}_2 \times 1.5 \text{ ATP} = 1.5 \text{ ATP}$$

$$1 \text{ GTP} \times 1 = 1 \text{ ATP}$$


$$\boxed{10 \text{ ATP}}$$





## Amphibolic nature of the citric acid cycle

Cataplerotic: out  
Anaplerotic: in



- Anaplerotic Rx<sup>n</sup>  
Synthesis of TCA cycle intermediate  
ATP =  $\downarrow$ , NADH =  $\downarrow$
- Cataplerotic Rx<sup>n</sup>  
• Decrease of TCA cycle intermediate  
• ATP =  $\uparrow$ , NADH =  $\uparrow$



## Gluconeogenesis

**Purpose:** Synthesis of glucose from non-carbohydrate precursors.

**Precursor:** Lactate, Pyruvate, Amino acids, Glycerol, Pyrimidine, Propionyl coA

**Pathway:** The reversal of glycolysis bypassing irreversible steps.


**Site:** The liver and kidney (mitochondria, Cytosol and ER), intestinal cells

**Condition:** Blood glucose level low, glucagon

**Regulation:** Glycolysis and gluconeogenesis are reciprocally regulated


cellular condition :  $ATP = \uparrow$      $Acetyl\ CoA = \uparrow$   
 $NADH = \uparrow$      $Urate = \uparrow$

Animals cannot convert acetyl coA into glucose.







Stage I: Conversion of Pyruvate to PEP




Pyruvate



(Biotin)  
Pyruvate carboxylase




Oxaloacetate



✗ → Citrate



Malate Dehydrogenase



malate


move out to cytosol



4C  
OAA

3C  
PEP

PEP carboxy Kinase



mitochondria = ATP = ↑  
acetyl CoA = ↑  
Citrate = ↑





Reversible

Stage 2: Conversion of PEP to FI6BP

Stage 3: Conversion of FI6BP to G6P

Fructose 16 Bis phosphate

X (IP) F16 BIS Phosphatase (active) — (P)

Fructose - 6 - P


isomerase

Glucose - 6 - P

Glucose

Glycolysis

Gluconeogenesis






## Stage 4: Conversion of G6P to Glucose

: Glucose-6-phosphatase

→ SER of liver & kidney.

Glucagon → → → → (P) (Active)

✓ GLUT-2 : Liver  
GLUT 4 : muscle




2 Pyruvate → 1 Glucose

- 4 ATP
  - 2 GTP
  - 2 NADH
- } Investment



### ✓ Cori-Cycle:

Lactate produced from glucose under anaerobic conditions in muscles and RBC is transported to liver converted into pyruvate then into glucose





## **PENTOSE PHOSPHATE PATHWAY or HEXOSE MONOPHOSPHATE SHUNT**

**Alternative pathway to glycolysis for the oxidation of glucose.**

Location: Cytosol

Provides: NADPH and Ribose-5-phosphate




## Purpose of Pentose Phosphate Pathway

Also provide pentose phosphate sugar

Nucleotide biosynthesis

NADPH: Anti-oxidant machinery of RBC



- Oxidize
- membrane lipids of RBC
- Hemolytic anemia

- Biosynthesis of aa, fatty acid, cholesterol nucleotides
- Detoxification rxn by cyt P 450 (SER of liver)
- Phagocytotic cells  $\rightarrow$  oxygen dependent killing



Consists of three stages – **Oxidation, Isomerization and Regeneration** - in which NADPH is produced, pentose undergo isomerization, and glycolytic intermediates are recovered.

NADP


NADPH

various pentose  
phosphate sugar




### Stage I: Oxidative phase:

Step 1: Glucose 6 Phosphate dehydrogenase  $\xrightarrow{NADP^+}$   $NADPH$




Step 2 & 3: 6-Phosphogluconolactonase and 6-Phosphogluconate dehydrogenase








## Stage 2: Isomerization and Epimerization of Ribulose-5-Phosphate





## Stage 3: Regeneration of Glycolytic intermediates involves Carbon-Carbon Bond Cleavage and Formation





## Glucose-6-Phosphate Dehydrogenase Deficiency

x-linked  
Recessive disorder

- NO NADPH production
- $H_2O_2 = \uparrow$
- Hemolytic anemia

- cannot consume → vicia faba  
favism
- Sensitive to anti malarial  
drugs.

Glutathione peroxidase, reductase  
Favabeans (vicia faba), primaquine



## Glycogen Biosynthesis

**Location:** Liver and Muscles (Cytosol)




### Physiological Conditions

- High Blood Glucose ✓
- Insulin Signaling ✓
- ATP High ✓
- NADH High ✓
- Glucose 6 Phosphate high ✓

**Active form:** UDP Glucose


## I. Synthesis of active glucosyl unit: UDP-Glucose

Liver  
Muscle





## 2. Glycogen extension by glycogen synthase : form $\alpha$ 1 $\rightarrow$ 4 Glycosidic bond.




-ve = ADP and Pi, Phosphorylated form (Glucagon/Epinephrine)  
+ = G6P, Dephosphorylated form (Insulin)




### 3. Glycogenin Primes Glycogen Synthesis

- 349-residue protein
- Acts as a glycosyltransferase
- Attaches a glucose residue donated by UDP-Glucose to the OH group of its Tyr 194
- Extends the glucose chain by up to seven additional UDP Glucose-donated glucose





## 4. Branching enzyme transfers seven-residue glycogen segments



$\alpha 1 \rightarrow 4$


Branch :  $\alpha 1 \rightarrow 6$  bond

Branching enzyme

Break  $\alpha 1 \rightarrow 4$  bond & form  $\alpha 1 \rightarrow 6$  bond.



The overall reaction of the glycogen synthesis for the addition of each glucose residue is





## Glycogen Breakdown (Glycogenolysis)

Conversions from glycogen to Glucose-1-P to Glucose-6-P and finally to Glucose.

### Physiological Conditions

Liver: Blood glucose = Low  
Glucagon signalling

cellular  
condition = ATP  $\uparrow$   
= NADH  $\uparrow$

Muscle: contraction  
ATP =  $\downarrow$ , NADH =  $\downarrow$



# I. Glycogen phosphorylase






## De-branching Enzyme – Single Bifunctional Enzyme


- ✓  $(\alpha 1 \rightarrow 4)$  Trans-glycosylase (glycosyl-transferase)
- ✓  $(\alpha 1 \rightarrow 6)$ - Glucosidase

9 4-I-P  
 ① Glucose






## Glucose-6-Phosphatase Generates Glucose in the Liver.





## Glucagon/Epinephrine Signaling

← Blood glucose = ↓



Glycogen

Glucose-1-P





# Thank you

Learn From India's Best Educators  
India's **No.1** EdTech Company for Graduates  
& Post Graduates Examination.

➡ Download **IFAS** App Now



ANDROID



IOS



WINDOWS





## Harshada Sharma

# Earned: ₹25,500

"As is widely recognized, IFAS stands as India's leading institute for CSIR NET examination preparation. My Personal philosophy has always revolved around helping others and I've had the pleasure of introducing the IFAS Online Course to 12 of my classmates and juniors, which in turn allowed me to earn 25,500 rupees. You too can join in and refer your friends, allowing you to earn a 5% rewards based on the fees each of your referred friends Pay"



### Step – 01 Invite Friends

Share your referral code or Link with your friends.

### Step – 02 Earn Cashback

On Every Purchase made using your referral code, you earn 5% instant cashback in your bank account.

### Step – 03 Refer More, Earn More

Keep referring as many friends as you can and keep **Earning** cash as much as you can



Friendship is Sharing, Caring, Giving & Helping Each Other...!